



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$m_1 r^{-m-1} \{ -nd + [r - \frac{2}{3}(m+1)r^{-1}d^2](\cos\alpha + \cos 2\alpha + \cos 3\alpha + \dots \cos n\alpha) \\ + (m+1)d(\cos^2\alpha + \cos^2 2\alpha + \cos^2 3\alpha + \dots \cos^2 n\alpha) \\ + \frac{1}{2}[(m+1)(m+3)]r^{-1}d^2(\cos^3\alpha + \cos^3 2\alpha + \cos^3 3\alpha + \dots \cos^3 n\alpha) \}.$$

By trigonometry,

$$\cos\alpha + \cos 2\alpha + \dots \cos n\alpha = \frac{\cos \frac{1}{2}(n+1)\alpha \cdot \sin \frac{1}{2}n\alpha}{\sin \frac{1}{2}\alpha};$$

$$\cos^2\alpha + \cos^2 2\alpha + \dots \cos^2 n\alpha = \frac{1}{2} \left\{ n + \frac{\cos(n+1)\alpha \cdot \sin n\alpha}{\sin \alpha} \right\};$$

$$\cos^3\alpha + \cos^3 2\alpha + \dots \cos^3 n\alpha = \frac{\cos \frac{1}{2}(n+3)\alpha \cdot \sin \frac{3}{2}n\alpha}{4\sin \frac{3}{2}\alpha} + \frac{3\cos \frac{1}{2}(n+1)\alpha \cdot \sin \frac{1}{2}n\alpha}{4\sin \frac{1}{2}\alpha}.$$

The value of these series when $n\alpha = 2\pi$ are 0, $n/2$, and 0, respectively.

Hence the expression for the approximate value of the resultant force reduces to

$$m_1 r^{-m-1} [-nd + (\frac{1}{2}n)(m+1)d], \text{ or } \frac{m_1 n(m-1)}{2r^{m+1}} d.$$

Also solved by G. B. M. ZERE.

PROBLEMS FOR SOLUTION.

ARITHMETIC.

95. Proposed by WALTER HUGH DRANE, A. M., Professor of Mathematics, Jefferson Military College, Washington, Miss.

Solve by arithmetic, if possible.

A man sold a house for \$7500 and gained a certain per cent. on the cost. If the cost had been 16 $\frac{2}{3}$ % less, his gain would have been 25% greater. Find the cost of the house.

96. Proposed by RAYMOND SMITH, Tiffin, Ohio.

How many acres in a square field whose diagonal is 10 rods longer than the side?

*** Solutions of these problems should be sent to B. F. Finkel, not later than April 10.

ALGEBRA.

85. Proposed by J. M. COLAW, A. M., Monterey, Va.

Sum the infinite series

$$\frac{1}{1^2 \cdot 2^2 \cdot 3^2 \cdot 4^2 \cdot 5^2} + \frac{1}{2^2 \cdot 3^2 \cdot 4^2 \cdot 5^2 \cdot 6^2} + \frac{1}{3^2 \cdot 4^2 \cdot 5^2 \cdot 6^2 \cdot 7^2} +, \text{ etc.}$$

86. Proposed by J. MARCUS BOORMAN, Consultative Mechanician, Counselor at Law, Inventor, Etc., Woodmere, Long Island, N. Y.

Solve $x^2 + yz = 16 \dots (A)$; $y^2 + xz = 17 \dots (B)$; $z^2 + xy = 22 \dots (C)$, for all the roots.

[This is Col. Titus' problem—see "Maseres' Tracts," pages 188-276—and is solved by Dr. Wallis in 51 pages, and by Mr. Frend in 38 pages, 8vo., but by the writer in 1 or 2 pages, 4to., or less. J. M. B.]

87. Proposed by the late B. F. BURLESON, Oneida Castle, N. Y.

A starts to travel around a circular island at a given point and travels at the rate of 5 miles in 4 hours. One half hour after A , B starts from a point directly opposite from A and travels in an opposite direction at the rate of 4 miles in 3 hours. One hour afterwards C starts from the same point as A and travels in an opposite direction to A at the rate of 3 miles in 2 hours. One half hour afterwards D starts from the same point as B and travels in an opposite direction to B at the rate of 2 miles in 1 hour. Required the size of the island, and when they will all be together, and how far each will have traveled at the accomplishment of this event.

*** Solutions of these problems should be sent to J. M. Colaw, not later than June 10.

GEOMETRY.

94. Proposed by EDMOND FISH, Hillsboro, Ill.

A tower $AB=a$, is surmounted by a flag pole $BC=b$. A point D is so taken in a line perpendicular to the foot of the tower that angle BDC is a maximum. Prove that AD is a mean proportional between AC and AB .

95. Proposed by WILLIAM HOOVER, A. M., Ph. D., Professor of Mathematics and Astronomy in Ohio State University, Athens, Ohio.

At each point of a parabola is described the rectangular hyperbola of four-pointic contact; prove that the locus of the center of the hyperbola is an equal parabola.

96. Proposed by W. F. BRADBURY, A. M., Head Master, Cambridge Latin School, Cambridge, Mass.

Isosceles triangles are constructed externally on the three sides of a triangle as bases, with the angles at the bases each 30° . The triangle formed by joining the remote vertices (the 120° vertices) of these isosceles triangles is equilateral. [Geometric—not Trigonometric—solution.]

*** Solutions of these problems should be sent to B. F. Finkel, not later than June 10.

CALCULUS.

75. Proposed by O. W. ANTHONY, M. Sc., Instructor in Mathematics in Boys' High School, New York City.

Solve the differential equation

$$\frac{d^2y}{dx^2} + n^2y = \frac{6}{x^2}y.$$

76. Proposed by E. B. ESCOTT, Cambridge, Mass.

Solve the partial differential equation

$$q^2r + 4pq s + p^2t + p^2q^2(rt - s^2) = a^2.$$

[Forsyth's *Differential Equations*, page 376.]

77. Proposed by T. E. COLE, Columbus, Ohio.

Derive the equation of a point in the pedal of a bicycle as the wheel rolls along on a plane.

*** Solutions of these problems should be sent to J. M. Colaw, not later than June 10.